

1800 SW 152nd Street, Ste 201, Burien, WA 98166 | (206) 248-1339 | www.kennerdentalgroup.com

DERMAL FILLER INFORMED CONSENT

in the following area(s):	;	
The indicated dermal fi	iller has been FDA approved for use in cosmetic tr	

I, ______understand that I will be injected with _____ dermal filler,

severe wrinkles around the nose and mouth. I understand this treatment is temporary, and reinjection is necessary after about six months. It has been explained to me that other temporary and more permanent treatments are available.

The following complications may occur with the dermal filler injection procedure:

- Risks: I understand there is a risk of bruising, redness, swelling, pain at the injection site, tenderness, itching, allergic reaction, and raised bumps of skin (nodules). These symptoms are usually mild and typically last a few days but can last up to a few months. In rare cases bruising can last several months and even be permanent.
- **2. Infection:** Post treatment bacterial, viral and/or fungal infections can occur which in most cases are easily treatable, however in rare cases permanent scarring in the area can occur.
- **3. Effectiveness:** Treatments can last anywhere from 4-6 months up to one year.
- **4. Treatments:** I understand more than one injection may be needed to achieve a satisfactory result.
- **5.** Allergic Reactions: In rare cases, there may be an allergic reaction to the injection.
- 6. There is a risk of scarring.
- 7. I will follow all aftercare instructions as it is crucial I do so for healing.

As dermal fillers are not an exact science, there might be an uneven appearance of the face with some areas more affected by the fillers than others. In most cases this uneven appearance can be corrected by more injections in the same or nearby areas. However in some cases this uneven appearance can persist for several weeks or months.

This list is not meant to be inclusive of all possible risks associated with dermal fillers as there are both known and unknown side effects associated with any medication or procedure.

These dermal fillers should not be administered to a pregnant or nursing woman.

The number of units injected is an estimate of the amount of dermal filler required to add volume to the skin and give the appearance of a smoother face. I understand there is no guarantee of results of any treatment and the regular charge applies to all subsequent treatments.

Patient Signature	Date:	
Witness_	Date:	